

ISSN 1027-4634

МАТЕМАТИЧНІ СТУДІЇ

MATEMATYCHNI
STUDII

Том 52, № 2 2019 Vol. 52, No. 2

МАТЕМАТИЧНІ СТУДІЇ

Праці Львівського математичного товариства

ISSN 2411-0620 (електронне видання)

ISSN 1027-4634 (друковане видання)

Журнал присвячено дослідженням в усіх областях математики. Приймаються оригінальні статті середнього розміру; виняток можливий для оглядових статей. Статті публікуються англійською, німецькою, російською, українською та французькою мовами. Виходить щоквартально.

Видається видавництвом «ВНТЛ-Класика» для Львівського математичного товариства.

Редактори:

М.М.Зарічний (Львів), О.Б.Скасків (Львів),
М.М.Шеремета (Львів)

Редакційна колегія:

О.Д. Артемович (Івано-Франківськ; Krakiv, Польща), Т.О.Банах (Львів), М.М.Бокало (Львів), Ю.Д.Головатий (Львів), Р.І.Григорчук (Москва, Росія), Р.О.Гринів (Львів), І.Й.Гуран (Львів), А.А.Дороговцев (Київ), Ю.А.Дрозд (Київ), Б.В.Забавський (Львів), С.Д.Івасишен (Київ), С.М.Івашкович (Лілль, Франція), В.В.Кириченко (Київ), Б.І.Копитко (Львів), А.А.Кореновський (Одеса), О.В.Лопушанський (Львів; Жешув, Польща), Я.В.Микитюк (Львів), Й.В.Островський (Харків; Ан卡拉, Туреччина), О.А.Панков (Балтімор, США), М.М.Попов (Чернівці), І.В.Протасов (Київ), О.Рашковський (Ставанґер, Норвегія), Д.Реповш (Любліана, Словенія), В.Г.Самойленко (Київ), А.М.Седлецький (Москва, Росія), С.Степін (Москва, Росія; Бялишток, Польща), О.Г.Сторож (Львів), С.Ю.Фаворов (Харків), Б.Н.Хабібулін (Уфа, Росія), І.Е.Чижиков (Львів)

Технічний редактор:

Д.Ю.Зікраб, А.О.Куриляк, А.І. Бандура

Адреса редакції:

Україна, 79000 Львів,
Університетська, 1
Львівський національний університет,
механіко-математичний факультет,
Математичні студії
e-mail: math.studii@lnu.edu.ua,
Вебсторінка: www.matstud.org.ua

Інформація про передплату:

З питань передплати звертатися за адресою:
ВНТЛ-Класика
а/с 10249, Львів, 79006
e-mail: info@vntl.com, http://www.vntl.com

MATEMATYCHNI STUDII

Proceedings of the Lviv Mathematical Society

ISSN 2411-0620 (Online)

ISSN 1027-4634 (Print)

Journal is devoted to research in all fields of mathematics. Original papers of moderate length are accepted; exception is possible for survey articles. Languages accepted are: English, German, Russian, and Ukrainian. Published quarterly.

Published for the Lviv Mathematical Society by VNTL Publishers.

Editors-in-Chief:

М.М.Sheremeta (Lviv), О.В.Skaskiv (Lviv),
М.М.Zarichnyi (Lviv)

Editorial Board:

O.D.Artemovych (Ivano-Frankivsk; Krakow, Poland), T.O.Banakh (Lviv), M.M.Bokalo (Lviv), I.E.Chyzhykov (Lviv), A.A.Dorogovtsev (Kyiv), Yu.A.Drozd (Kyiv), S.Yu.Favorov (Kharkiv), Yu.D.Golovaty (Lviv), R.I.Grigorchuk (Moscow,Russia), I.Yo.Guran (Lviv), R.O. Hrynniv (Lviv), S.D.Ivasyshen (Kyiv), S.M.Ivashkovych (Lille, France), B.N.Khabibullin (Ufa,Russia), V.V.Kirichenko (Kyiv), B.I.Kopitko (Lviv), A.A.Korenovskii (Odesa), O.V.Lopushanskyi (Lviv; Rzeszów,Poland), Ya.V.Mykytyuk (Lviv), I.V.Ostrovskii (Kharkiv; Ankara, Turkey), O.A. Pankov (Baltimore, USA), M.M.Popov (Chernivtsi), I.V.Protasov (Kyiv), D.Repovš (Ljubljana, Slovenia), V.G.Samoilenko (Kyiv), A.M.Sedletskii (Moscow, Russia), S.Stepin (Moscow, Russia; Bialystok, Poland), O.G.Storozh (Lviv), B.V.Zabavskyi (Lviv)

Technical editors:

A.O.Kuryliak, D.Yu.Zikrach, A.I. Bandura

Address:

Matematychni Studii
Dept. of Mechanics and Mathematics,
Lviv National University,
1 Universytetska St.,
79000, Lviv, Ukraine,
e-mail: math.studii@lnu.edu.ua
Web-site: www.matstud.org.ua

Subscription information:

Subscription information can be obtained by the address:
VNTL publishers
P.O.Box 10249, Lviv, 79006, Ukraine
e-mail: info@vntl.com, http://www.vntl.com

ПРАВИЛА ДЛЯ АВТОРІВ

1. При оформленні варто орієнтуватися на правила, прийняті в більшості математичних журналів (вони істотно не відрізняються). Оформлення списку літератури можна запозичити з даного випуску “Математичних студій”. Статті приймаються англійською, німецькою, російською, українською та французькою мовами.

2. Кожен рукопис повинен містити: (i) прізвище, ініціали автора, резюме англійською мовою (не більше ніж 250 слів, бажано без формул і цитування літератури); (ii) індекс УДК; (iii) 2010 Mathematics Subject Classification; (iv) якщо заголовок довгий, то подати також його короткий варіант, не більше 40 знаків; (v) адреса автора для публікації в журналі.

3. У випадку авторського колективу вказати прізвище того з авторів, з ким редколегія може вести листування.

4. З 1999 року введено розділ ”Короткі повідомлення”. Крім тексту коротких повідомлень (до 5 журнальних сторінок), автори зобов'язані прислати повні доведення всіх результатів.

5. Обов'язкове використання L^AT_EX' у при наборі статті за допомогою стилювого та макетного файлів журналу „Математичні студії“. Редколегія розглядає *лише* праці, які надіслані через систему автоматичної подачі статей на сторінці <http://matstud.org.ua>

INSTRUCTIONS TO AUTHORS

1. The conventions of mathematical journals should be observed. The style of references should be as in the papers published in the current issue. Languages accepted are: English, French, German, Russian, and Ukrainian.

2. Each typescript should include: (i) the author's name, title and an abstract of not more than 250 words written in English, preferably containing no formulae and certainly no complicated formulae and no references; (ii) a 2010 Mathematics Subject Classification; (iii) if the title is long, a shortened form of it, no more than forty characters in length; (iv) the address of the author(s) for publication in the journal.

3. In the case of joint authorship, a letter should accompany the manuscript stating which author will assume responsibility for correspondence and proofreading.

4. Since 1999, the section “Research Announcements” was introduced. Besides the text (up to 5 journal pages), the authors should submit an expanded version containing complete proofs of all results.

5. The papers should be typed in L^AT_EX using the style and maket files of the journal “Matematychni Studii”. The editorial board considers *only* the papers which are submitted by online submission system on the web-site <http://matstud.org.ua>

3 M I C T

BILANYK I. B. A truncation error bound for some branched continued fractions of a special form	115
NAKONECHNIY O. G., KUDIN G. I., ZINKO T. P. Formulas of perturbation for one class of pseudo inverse operators	124
BILOUS A. M. Idempotent matrices over a duo ring	133
SHEREMETA M. M., TRUKHAN YU. S. Properties of analytic solutions of a differential equation	138
MALYUTIN K. G., REVENKO A. A. Extremal problems in the space of meromorphic functions of finite order in the half-plane.....	144
ZABOLOTSKYI M. V., BASIUK YU. V., MOSTOVA M. R. Binomial asymptotics for the logarithmic derivative of zero-order entire functions with zeros along curves of regular rotation	156
MOKHON'KO A. Z., KOLYASA L. I. Some properties of meromorphic solutions of linear differential equation with meromorphic coefficients	166
EREMENKO A. A problem of Stanisław Saks	173
BAK S. M. Homoclinic traveling waves in discrete sine–Gordon equation with nonlinear interaction on 2D-lattice.....	176
SALIM D., BUDHI W. S., SOEHARYADI Y. On Rough maximal inequalities: an extension of Fefferman–Stein results	185
AMMAR A., JERIBI A., SAADAOUI B. On some classes of demicompact linear relation and some results of essential pseudospectra.....	195
BOUCHELAGHEM F., BENHARRAT M. On the spectrum of linear operator pencils	211
АВТОРСЬКИЙ ВКАЗІВНИК, 2019	222

C O N T E N T S

BILANYK I. B. A truncation error bound for some branched continued fractions of a special form	115
NAKONECHNIY O. G., KUDIN G. I., ZINKO T. P. Formulas of perturbation for one class of pseudo inverse operators	124
BILOUS A. M. Idempotent matrices over a duo ring.....	133
SHEREMETA M. M., TRUKHAN YU. S. Properties of analytic solutions of a differential equation	138
MALYUTIN K. G., REVENKO A. A. Extremal problems in the space of meromorphic functions of finite order in the half-plane.....	144
ZABOLOTSKYI M. V., BASIUK YU. V., MOSTOVA M. R. Binomial asymptotics for the logarithmic derivative of zero-order entire functions with zeros along curves of regular rotation	156
MOKHON'KO A. Z., KOLYASA L. I. Some properties of meromorphic solutions of linear differential equation with meromorphic coefficients.....	166
EREMENKO A. A problem of Stanisław Saks	173
BAK S. M. Homoclinic traveling waves in discrete sine–Gordon equation with nonlinear interaction on 2D-lattice.....	176
SALIM D., BUDHI W. S., SOEHARYADI Y. On Rough maximal inequalities: an extension of Fefferman–Stein results	185
AMMAR A., JERIBI A., SAADAQUI B. On some classes of demicompact linear relation and some results of essential pseudospectra	195
BOUCHELAGHEM F., BENHARRAT M. On the spectrum of linear operator pencils	211
INDEX, 2019	222

I. B. BILANYK

A TRUNCATION ERROR BOUND FOR SOME BRANCHED CONTINUED FRACTIONS OF A SPECIAL FORM

I. B. Bilanyk. *A truncation error bound for some branched continued fractions of a special form*, Mat. Stud. **52** (2019), 115–123.

By analogy with continued fractions, the angular domain of convergence of branched continued fraction of a special form is established. The truncation error bound for branched continued fractions of the special form is established. For this we have used the estimates for continued fractions obtained by J. Jensen, W. Gragg, D. Warner, the multidimensional analogue of the van Vleck Theorem, the analytic theory of continued fractions and branched continued fractions and elements of the theory of stability under perturbations. In comparison with already known results, in the paper, isolation conditions from zero for elements of branched continued fractions of the special form are weakened, but there are some requirements for the elements' rate of tending to zero. The obtained result is a multidimensional analogue of estimates for van Vleck continued fractions.

Introduction. One of the classical convergence criteria of continued fractions is the van Vleck Theorem [18, 19], which contains sufficient conditions of convergence of continued fractions with complex elements that lie in the angular domain. The proof is based on using the Stieltjes–Vitaly Theorem and it does not need obtaining truncation error bounds.

In the theory of continued fractions, there are some theorems where the fact of convergence is proved but the estimate of approximation is not. For example, it has not known yet the truncation error bound for continued fractions in the case when the elements satisfy the conditions of the Sleszynski–Pringsheim Theorem [5].

J. Jensen [14, 18] proved the estimates of convergence rate for van Vleck continued fractions. Another method to prove a similar estimate is showed by W. Gragg and D. Warner in the paper [13]. The method from that paper was also used for establishing the truncation error bounds for continued fractions in parabolic domains.

Theorem 1 (W. Gragg, D. Warner, [13]). *Let the elements of the continued fraction*

$$b_0 + \cfrac{1}{b_1 + \cfrac{1}{b_2 + \dots}} = b_0 + \overline{\overline{D}}_{n=1}^{\infty} \cfrac{1}{b_n}$$

satisfy the conditions

$$b_k \neq 0, \quad |\arg b_k| < \theta, \quad \theta < \frac{\pi}{2}, \quad k = 0, 1, 2, \dots . \quad (1)$$

2010 Mathematics Subject Classification: 11J70, 30B70, 31A55, 40A15.

Keywords: continued fraction; branched continued fraction of the special form; truncation error bound.
doi:10.30970/ms.52.2.115-123

Then

- (i) there exist the finite limits of even and odd approximants;
- (ii) the sequence of approximants $\{f_n\}$ converges iff the series $\sum_{n=1}^{\infty} |b_n|$ diverges;
- (iii) the bound

$$|f_m - f_{n-1}| \leq \frac{1}{d_n}, \quad m \geq n, \quad (2)$$

holds with

$$d_n \geq \frac{\operatorname{Re} b_1}{2 + \operatorname{Re} b_1} \cos \theta \ln \left(1 + (\operatorname{Re}(b_1))^2 \min \left\{ 1, \frac{1}{|b_1|^2} \right\} \cos \theta \sum_{k=1}^n |b_k| \right), \quad n \geq 1.$$

An analogue of the van Vleck Theorem is established for branched continued fractions (BCF) [5], two-dimensional continued fractions [10, 16], BCF of a special form [8]. By T. Antonova in the paper [1] the truncation error bound in an angular domain for BCF of the special form is obtained in case when elements are isolated from zero. The bound for two-dimensional continued fractions in an angular domain with putting on additional conditions is proved by O. Sus' [17]. These conditions are requirements regarding the speed of divergence to zero of certain product that is constructed of elements of the two-dimensional continued fraction. In [8] there is a truncation error bound for BCF of the special form with elements isolated from zero. This result is similar to J. Jensen estimate.

In the paper, the truncation error bound is established for BCF of the special form

$$b_0 + \overline{D} \sum_{k=1}^{\infty} \sum_{i_k=1}^{i_{k-1}} \frac{1}{b_{i(k)}} = b_0 + \sum_{i_1=1}^N \frac{1}{b_{i(1)} + \sum_{i_2=1}^{i_1} \frac{1}{b_{i(2)} + \sum_{i_3=1}^{i_2} \frac{1}{b_{i(3)} + \dots}}}, \quad (3)$$

where $b_0, b_{i(k)} \in \mathbb{C}$, $i(k) \in \mathcal{I}$,

$$\mathcal{I} = \{i(k) = (i_1, i_2, \dots, i_k) : 1 \leq i_k \leq i_{k-1} \leq \dots \leq i_0; k \geq 1; i_0 = N\},$$

N is a fixed natural number that determines the dimension of (3).

The n th approximant of (3) is the expression

$$f_n = b_0 + \overline{D} \sum_{k=1}^n \sum_{i_k=1}^{i_{k-1}} \frac{1}{b_{i(k)}}.$$

The BCF (3) converges if there exists the finite limit $\lim_{n \rightarrow \infty} f_n = f$. The number f is called the value of (3).

Tails of n th approximant of (3) are following finite BCF

$$Q_{i(n)}^{(n)} = b_{i(n)}, \quad Q_{i(p)}^{(n)} = b_{i(p)} + \overline{D} \sum_{r=p+1}^n \sum_{i_r=1}^{i_{r-1}} \frac{1}{b_{i(r)}}, \quad (4)$$

where $n = 1, 2, \dots$; $p = \overline{1, n-1}$; $i(n) \in \mathcal{I}$; $i(p) \in \mathcal{I}$.

The convergence problem and researching truncation error bounds of BCF of the special form in different domains are considered in a large number of papers, in particular, by D. Bodnar [6, 9], O. Baran [3, 4], T. Antonova [1, 2], R. Dmytryshyn [11, 12], etc.

Henceforward we will use the following formulas that gives a possibility to estimate absolute errors of calculating a finite continued fraction.

Let

$$f_n = b_0 + \overline{\sum_{k=1}^n \frac{1}{b_k}}, \quad \tilde{f}_n = \tilde{b}_0 + \overline{\sum_{k=1}^n \frac{1}{\tilde{b}_k}}, \quad b_k, \tilde{b}_k \in \mathbb{C}, \quad k = 0, 1, \dots, n; \quad n \geq 1.$$

Then ([7, 15])

$$f_n - \tilde{f}_n = b_0 - \tilde{b}_0 + (-1)^n \sum_{k=1}^n \frac{b_k - \tilde{b}_k}{\prod_{r=1}^k Q_r^{(n)} \tilde{Q}_r^{(n)}}, \quad (5)$$

where

$$Q_r^{(n)} = b_r + \overline{\sum_{m=r+1}^n \frac{1}{b_m}}, \quad \tilde{Q}_r^{(n)} = \tilde{b}_r + \overline{\sum_{m=r+1}^n \frac{1}{\tilde{b}_m}}$$

are the tails of the corresponding finite continued fractions.

Main result. We consider the subsets $\mathcal{I}^{(m)}$ of the set of multiindices \mathcal{I} ,

$$\mathcal{I}^{(m)} = \{i(n) = (i_1, i_2, \dots, i_n) : m \leq i_n \leq i_{n-1} \leq \dots \leq i_0; \quad n \geq 1; \quad i_0 = N\}, \quad m = \overline{2, N},$$

and we set

$$s[k] = (\underbrace{s, s, \dots, s}_k), \quad 1 \leq s \leq N, \quad k = 1, 2, \dots$$

The main result of the paper is the following theorem that is an analog of Theorem 1 for BCF of the special form.

Theorem 2. Let there exist constants θ, δ, β such that $0 < \theta < \frac{\pi}{4}$, $0 < \delta < 1$, $0 \leq \beta \leq \frac{1}{2}$ and the elements of the BCF of the special form (3) satisfy the conditions

$$|\arg b_0| \leq \theta, \quad |\arg b_{i(k)}| \leq \theta, \quad i(k) \in \mathcal{I}, \quad (6)$$

$$\operatorname{Re}(b_0) \geq \delta, \quad \operatorname{Re}(b_{i(n)}) \geq \delta, \quad \operatorname{Re}(b_{1[s]}) \geq \frac{\delta}{s^\beta}, \quad \operatorname{Re}(b_{i(n)1[s]}) \geq \frac{\delta}{s^\beta}, \quad s \geq 1, \quad i(n) \in \mathcal{I}^{(2)}. \quad (7)$$

Then BCF (3) converges and the following truncation error bound holds

$$|f_m - f_{Nn}| < \frac{M_N}{\ln \left(1 + \frac{\alpha}{1-\beta} \left((n+1)^{1-\beta} - 1 \right) \right)}, \quad m \geq Nn, \quad n \in \mathbb{N}, \quad (8)$$

where M_N and α are positive numbers that do not depend on n and m ,

$$\begin{aligned} M_N &= \frac{2+\delta}{\delta \cos \theta} K_N, \quad K_1 = 1, \quad K_j = 4AK_{j-1} + 2, \quad 2 \leq j \leq N, \\ A &= \left(1 + \frac{1}{\delta^2} \right) \left(1 + \frac{1}{\delta^4 \cos^2 2\theta + 2\delta^2 \cos 2\theta} \right). \end{aligned} \quad (9)$$

Proof. The analogue of the van Vleck Theorem for BCF of the special form [8] proves that the BCF (3) converges.

We shall prove estimate (8) by mathematical induction in N , the dimension of the BCF of the special form (3).

If $N = 1$ the BCF (3) becomes the continued fraction

$$b_0 + \frac{1}{b_1 + \frac{1}{b_{11} + \frac{1}{b_{111} + \dots}}}. \quad (10)$$

Let f_k be its k th approximant, $k \geq 0$. From condition (6) it follows that the assumptions of Theorem 1 for the continued fraction (10) hold. In consequence, we obtain the estimate

$$|f_m - f_n| \leq \frac{1}{d_{n+1}} < \frac{1}{\mu_n}, \quad m \geq n,$$

where

$$\mu_n = \frac{\operatorname{Re} b_1}{2 + \operatorname{Re} b_1} \cos \theta \ln \left(1 + (\operatorname{Re} b_1)^2 \min \left\{ 1, \frac{1}{|b_1|^2} \right\} \cos \theta \sum_{k=1}^n |b_{1[k]}| \right), \quad n \geq 1.$$

It is obvious that

$$\frac{\operatorname{Re} b_1}{2 + \operatorname{Re} b_1} \geq \frac{\delta}{2 + \delta}.$$

Considering the conditions (6), (7) of Theorem 2, we get

$$(\operatorname{Re} b_1)^2 \min \left\{ 1, \frac{1}{|b_1|^2} \right\} = \min \{ (\operatorname{Re} b_1)^2, \cos^2(\arg b_1) \} \geq \min \{ \delta^2, \cos^2 \theta \}$$

and

$$\sum_{k=1}^n |b_{1[k]}| \geq \sum_{k=1}^n \frac{\delta}{k^\beta} \geq \delta \int_1^{n+1} \frac{dx}{x^\beta} = \delta \frac{(n+1)^{1-\beta} - 1}{1 - \beta},$$

because the function $y = \frac{1}{x^\beta}$ monotonically decreases in x , $x > 0$.

Thus,

$$|f_m - f_n| < \frac{2 + \delta}{\delta \cos \theta} \frac{1}{\ln \left(1 + \frac{\alpha}{1 - \beta} ((n+1)^{1-\beta} - 1) \right)}, \quad m \geq n,$$

where $\alpha = \min \{ \delta^3 \cos \theta, \delta \cos^3 \theta \}$, that is, the inequality (8) is true when $N = 1$ where $M_1 = \frac{2+\delta}{\delta \cos \theta}$.

Suppose that the estimate (8) holds for $N = r - 1$. Let us now prove it for $N = r$. Consider the r -dimensional BCF of the special form (3) where $i_0 = r$. Write down its n th approximant in the form

$$f_n = b_0^{(r-1,n)} + \overline{D}_{k=1}^n \frac{1}{b_{r[k]}^{(r-1,n-k)}}, \quad n \geq 1,$$

where

$$b_0^{(r-1,n)} = b_0 + \overline{D}_{l=1}^n \sum_{i_l=1}^{i_{l-1}} \frac{1}{b_{i(l)}}, \quad b_{r[n]}^{(r-1,0)} = b_{r[n]}, \quad b_{r[k]}^{(r-1,n-k)} = b_{r[k]} + \overline{D}_{l=1}^{n-k} \sum_{i_l=1}^{i_{l-1}} \frac{1}{b_{r[k]i(l)}}, \quad (11)$$

$k = 1, 2, \dots, n - 1$, $i_0 = r - 1$, are the approximants of respective $(r - 1)$ -dimensional BCF that are contained in the structure of BCF (3).

Let

$$\hat{f}_n = b_0^{(r-1)} + \overline{\text{D}}_{k=1}^n \frac{1}{b_{r[k]}^{(r-1)}}$$

are the n th approximant of continued fraction that is constructed of all calculated $(r - 1)$ -dimensional BCF in the BCF (3), that is,

$$b_0^{(r-1)} = b_0 + \overline{\text{D}}_{l=1}^{\infty} \sum_{i_l=1}^{i_{l-1}} \frac{1}{b_{i(l)}}, \quad b_{r[k]}^{(r-1)} = b_{r[k]} + \overline{\text{D}}_{l=1}^{\infty} \sum_{i_l=1}^{i_{l-1}} \frac{1}{b_{r[k]i(l)}}, \quad (12)$$

$$i_0 = r - 1, \quad k = 1, 2, \dots, n.$$

These BCF converge in accordance with the analogue of the van Vleck Convergence Theorem for BCF of the special form [8].

Then

$$|f_m - f_{rn}| \leq |f_m - \hat{f}_n| + |f_{rn} - \hat{f}_n|, \quad m \geq rn.$$

Let us now estimate the values $|f_p - \hat{f}_n|$, $p \geq rn$.

Consider the next finite continued fraction

$$h_{p,n} = b_0^{(r-1)} + \frac{1}{b_{r[1]}^{(r-1)}} + \dots + \frac{1}{b_{r[n]}^{(r-1)}} + \frac{1}{b_{r[n+1]}^{(r-1,p-n-1)}} + \dots + \frac{1}{b_{r[p]}^{(r-1,0)}},$$

with elements that are defined in accordance with (11), (12). Obviously,

$$|f_p - \hat{f}_n| \leq |f_p - h_{p,n}| + |h_{p,n} - \hat{f}_n|. \quad (13)$$

Let us now obtain estimations from above for each addend in the right-hand side of inequality (13). From the formula (5) it follows that

$$|f_p - h_{p,n}| \leq |b_0^{(r-1)} - b_0^{(r-1,p)}| + \sum_{k=1}^n \frac{|b_{r[k]}^{(r-1)} - b_{r[k]}^{(r-1,p-k)}|}{\prod_{s=1}^k |\tilde{Q}_{r[s]}^{(p)} Q_{r[s]}^{(p)}|}, \quad (14)$$

where $\tilde{Q}_{r[s]}^{(p)}$, $Q_{r[s]}^{(p)}$ are s th tails of the continued fractions $h_{p,n}$ and f_p , respectively, $s = \overline{1, k}$.

Consider the products in the denominators of the previous relation. If $k = 2l$, then

$$\prod_{s=1}^{2l} |\tilde{Q}_{r[s]}^{(p)} Q_{r[s]}^{(p)}| = \prod_{s=1}^l \left(|\tilde{Q}_{r[2s-1]}^{(p)} \tilde{Q}_{r[2s]}^{(p)}| \left| Q_{r[2s-1]}^{(p)} Q_{r[2s]}^{(p)} \right| \right).$$

Taking into account

$$\operatorname{Re}(b_{r[k]}^{(r-1)}) = \operatorname{Re}(b_{r[k]}) + \operatorname{Re} \left(\overline{\text{D}}_{l=1}^{\infty} \sum_{i_l=1}^{i_{l-1}} \frac{1}{b_{r[k]i(l)}} \right) > \operatorname{Re}(b_{r[k]}) \geq \delta, \quad k = \overline{1, n},$$

we estimate separately each factor of product

$$\begin{aligned} \left| \tilde{Q}_{r[2s-1]}^{(p)} \tilde{Q}_{r[2s]}^{(p)} \right| &= \left| \tilde{Q}_{r[2s]}^{(p)} \left(b_{r[2s-1]}^{(r-1)} + \frac{1}{\tilde{Q}_{r[2s]}^{(p)}} \right) \right| \geq \operatorname{Re} \left(b_{r[2s-1]}^{(r-1)} \tilde{Q}_{r[2s]}^{(p)} + 1 \right) = \\ &= \operatorname{Re} \left(b_{r[2s-1]}^{(r-1)} \left(b_{r[2s]}^{(r-1)} + \frac{1}{\tilde{Q}_{r[2s+1]}^{(p)}} \right) \right) + 1 > \operatorname{Re} (b_{r[2s-1]} b_{r[2s]}) + 1 \geq \delta^2 \cos 2\theta + 1. \end{aligned}$$

For factors $\left| Q_{r[2s-1]}^{(p)} Q_{r[2s]}^{(p)} \right|$ the analogous estimations are valid.

Therefore,

$$\prod_{s=1}^{2l} \left| \tilde{Q}_{r[s]}^{(p)} Q_{r[s]}^{(p)} \right| > (\delta^2 \cos 2\theta + 1)^{2l}.$$

If $k = 2l + 1$, then

$$\begin{aligned} \prod_{s=1}^{2l+1} \left| \tilde{Q}_{r[s]}^{(p)} Q_{r[s]}^{(p)} \right| &= \left| \tilde{Q}_{r[1]}^{(p)} Q_{r[1]}^{(p)} \right| \prod_{s=1}^l \left(\left| \tilde{Q}_{r[2s]}^{(p)} \tilde{Q}_{r[2s+1]}^{(p)} \right| \left| Q_{r[2s]}^{(p)} Q_{r[2s+1]}^{(p)} \right| \right) > \\ &> (\operatorname{Re} (b_{r[1]})) ^2 \prod_{s=1}^l (\operatorname{Re} (b_{r[2s]} b_{r[2s+1]}) + 1)^2 \geq \delta^2 (\delta^2 \cos 2\theta + 1)^{2l}. \end{aligned}$$

Hence,

$$\prod_{s=1}^k \left| \tilde{Q}_{r[s]}^{(p)} Q_{r[s]}^{(p)} \right| > \delta^{1-(-1)^k} (\delta^2 \cos 2\theta + 1)^{2[\frac{k}{2}]}, \quad 1 \leq k \leq n.$$

We shall estimate the values of numerators

$$\left| b_{r[k]}^{(r-1)} - b_{r[k]}^{(r-1,p-k)} \right|,$$

$1 \leq k \leq n$, of the inequality (14) as the difference between the value of an $(r-1)$ -dimensional BCF of the special form and its $(p-k)$ th approximant, $p-k \geq (r-1)n$. By the inductive hypothesis, taking into account Theorem 1, we get

$$\left| b_{r[k]}^{(r-1)} - b_{r[k]}^{(r-1,p-k)} \right| < \frac{2+\delta}{\delta \cos \theta} \frac{2K_{r-1}}{\ln \left(1 + \frac{\alpha}{1-\beta} ((n+1)^{1-\beta} - 1) \right)}, \quad k = 1, 2, \dots, n, \quad p \geq rn.$$

The analogous estimation is also true for the first addend of the right-hand side of the inequality (14).

In these terms,

$$\begin{aligned} |f_p - h_{p,n}| &< \frac{2+\delta}{\delta \cos \theta} \frac{2K_{r-1}}{\ln \left(1 + \frac{\alpha}{1-\beta} ((n+1)^{1-\beta} - 1) \right)} \\ &\left(1 + \frac{1}{\delta^2} + \frac{1}{(\delta^2 \cos 2\theta + 1)^2} + \frac{1}{\delta^2 (\delta^2 \cos 2\theta + 1)^2} + \dots + \frac{1}{\delta^{1-(-1)^n} (\delta^2 \cos 2\theta + 1)^{2[\frac{n}{2}]}} \right) < \end{aligned}$$

$$\begin{aligned}
&< \frac{2+\delta}{\delta \cos \theta} \frac{2K_{r-1}}{\ln \left(1 + \frac{\alpha}{1-\beta} ((n+1)^{1-\beta} - 1) \right)} \left(1 + \frac{1}{\delta^2} \right) \sum_{k=0}^{\infty} (\delta^2 \cos 2\theta + 1)^{-2k} = \\
&= \frac{2+\delta}{\delta \cos \theta} \frac{2K_{r-1}A}{\ln \left(1 + \frac{\alpha}{1-\beta} ((n+1)^{1-\beta} - 1) \right)}, \quad p \geq rn,
\end{aligned}$$

where A is defined according to the formula (9).

Let us estimate the second addend in the right-hand side inequality (13) $|h_{p,n} - \hat{f}_n|$, which is a difference of approximants of some continued fractions with elements from the domain (1). Similarly with (2) we obtain

$$|h_{p,n} - \hat{f}_n| < \frac{1}{\nu_n}, \quad n \geq 1,$$

where

$$\nu_n = \frac{\operatorname{Re} (b_r^{(r-1)})}{2 + \operatorname{Re} (b_r^{(r-1)})} \cos \theta \ln \left(1 + (\operatorname{Re} (b_r^{(r-1)}))^2 \min \left\{ 1, \frac{1}{|b_r^{(r-1)}|^2} \right\} \cos \theta \sum_{s=1}^n |b_{r[s]}^{(r-1)}| \right).$$

Since $\operatorname{Re} b_{r[k]}^{(r-1)} > \delta$, $k = 1, 2, \dots$, taking into account the conditions of Theorem 2, we have

$$\nu_n > \frac{\delta \cos \theta}{2 + \delta} \ln (1 + \alpha n), \quad n \geq 1.$$

Then

$$|h_{p,n} - \hat{f}_n| < \frac{2+\delta}{\delta \cos \theta} \frac{1}{\ln (1 + \alpha n)}.$$

Using the previous estimates of the addends in the right-hand side the equation (13), we obtain

$$|\hat{f}_p - \hat{f}_n| < \frac{2+\delta}{\delta \cos \theta} \frac{2K_{r-1}A + 1}{\ln \left(1 + \frac{\alpha}{1-\beta} ((n+1)^{1-\beta} - 1) \right)}, \quad p \geq rn,$$

because

$$n > \frac{1}{1-\beta} ((n+1)^{1-\beta} - 1).$$

For $m \geq rn$, putting in the last inequality $p = m$, $p = rn$, we get

$$\begin{aligned}
&|f_m - f_{rn}| \leq |f_m - \hat{f}_n| + |\hat{f}_{rn} - \hat{f}_n| < \\
&< \frac{2+\delta}{\delta \cos \theta} \frac{2(2K_{r-1}A + 1)}{\ln \left(1 + \frac{\alpha}{1-\beta} ((n+1)^{1-\beta} - 1) \right)} = \frac{M_r}{\ln \left(1 + \frac{\alpha}{1-\beta} ((n+1)^{1-\beta} - 1) \right)},
\end{aligned}$$

where

$$M_r = \frac{2+\delta}{\delta \cos \theta} K_r, \quad K_r = 2(2K_{r-1}A + 1).$$

Consequently, for approximants f_k of N -dimensional BCF (3), where N is an arbitrary fixed natural number, the following estimation is true

$$|f_m - f_{Nn}| < \frac{M_N}{\ln \left(1 + \frac{\alpha}{1-\beta} \left((n+1)^{1-\beta} - 1 \right) \right)}, \quad m \geq Nn,$$

where

$$M_N = \frac{2+\delta}{\delta \cos \theta} K_N, \quad N \geq 1, \quad K_1 = 1, \quad K_j = 4AK_{j-1} + 2, \quad 2 \leq j \leq N.$$

□

Conclusion. The obtained truncation error bound for some BCF of a special form in the domain that is narrower than its convergence domain. Thereby, the problem of establishing estimations in the whole convergence domain is still unsolved. The results of Theorem 2 can be used for establishing truncation error bounds for functional BCF with independent variables, in particular, multidimensional J -fractions with independent variables.

REFERENCES

1. T.M. Antonova, *Convergence rate of branched continued fractions of the special form*, Volyn Math. Bulletin, **6** (1999), 3–8. (in Ukrainian)
2. T.M. Antonova, D.I. Bodnar, Domains of convergence of branched continued fractions of a special form, in: Approximation Theory of Functions and Its Application: Proceedings of the Institute of Mathematics, V.31, Institute of Mathematics, Ukrainian National Academy of Sciences (2000), 19–32. (in Ukrainian)
3. O.E. Baran, *An analogues of the Vorpits'kii convergence criterion for branched continued fractions of the special form*, Mat. Met. Fiz.-Mekh. Polya, **39** (1996), №2, 35–38; English translation: J. Math. Sci., **90** (1998), №5, 2348–2351.
4. O.E. Baran, *Some domains of convergence of branched continued fractions of a special form*, Carpathian Math. Publ., **5**, №1 (2013), 4–13. (in Ukrainian)
5. D.I. Bodnar, Branched Continued Fractions, Naukova Dumka, Kyiv, 1986. (in Russian)
6. D.I. Bodnar, *Investigation of the convergence of a class of branching continued fractions*, Continued fractions and its application, Institute of Mathematics, Kyiv, 1976, 41–44. (in Russian)
7. D.I. Bodnar, I.B. Bilanyk, *Convergence criterion for branched continued fractions of the special form with positive elements*, Carpathian Math. Publ., **9** (2017), №1, 13–21.
8. D.I. Bodnar, I.B. Bilanyk, *On the convergence of branched continued fractions of the special form in angular domains*, Mat. Met. Fiz.-Mekh. Polya, **60** (2017), №3, 60–69.
9. D.I. Bodnar, R.I. Dmytryshyn, *On the convergence of multidimensional g-fraction*, Mat. Stud., **15** (2001), №2, 115–126.
10. D.I. Bodnar, Kh.Yo. Kuchminska, *An analog of the van Vleck theorem for two-dimensional continued fractions*, Mat. Met. Fiz.-Mekh. Polya, **42** (1999), №4, 21–25. (in Ukrainian)
11. O.S. Bodnar, R.I. Dmytryshyn, *On the convergence of multidimensional S-fractions with independent variables*, Carpathian Math. Publ., **10** (2018), №1, 58–64.
12. R.I. Dmytryshyn, *On the convergence criterion for branched continued fractions with independent variables*, Carpathian Math. Publ., **9** (2017), №2, 120–127.
13. W.B. Gragg, D.D. Warner, *Two constructive results in continued fractions*, SCIAM J. Numer. Anal., **20** (1983), №3, 1187–1197.
14. J.L. W.V. Jensen, Bidrag til Kaedebrekernes Teori. Festschrift til H.G. Zeuthen, 1909.
15. V.R. Hladun, Analysis of stability under perturbations of branched continued fractions, Candidate-Degree Thesis (Physics and Mathematics), Lviv, 2007. (in Ukrainian)

16. Kh.Yo. Kuchmins'ka, Two-Dimensional Continued Fractions, Pidstryhach Institute for Applied Problems in Mechanics and Mathematics, Ukrainian National Academy of Sciences, Lviv, 2010. (in Ukrainian)
17. O.M. Sus', *On the evaluation of the rate of convergence of two-dimensional continued fractions with complex elements*, Prykl. Probl. Mekh. Mat., **6** (2008), 115–123. (in Ukrainian).
18. O. Perron, Die Lehre von den Kettenbrüchen, Leipzig; Berlin: Druck und verlag von B.G. Teubner, 1913.
19. E.B. van Vleck, *On the convergence of continued fractions with complex elements*, Trans. Amer. Math. Soc., **2** (1901), 215–233.

Pidstryhach Institute for Applied Problems
of Mechanics and Mathematics, NASU
Lviv, Ukraine
i.bilanyk@ukr.net

Received 27.09.2019