

Wiadomości Lekarskie

Medical Advances

Official journal of the Polish Medical Association
Wiadomości Lekarskie has been published since 1928



Volume LXXVII, Issue 12, DECEMBER 2024

ISSN 0043-5147

E-ISSN 2719-342X

Wiadomości Lekarskie

Medical Advances

Official journal of the Polish Medical Association

Wiadomości Lekarskie has been published since 1928



Volume LXXVII, Issue 12, DECEMBER 2024



Memory of
dr Władysław
Biegański

Wiadomości Lekarskie Medical Advances is abstracted and indexed in:
PUBMED/MEDLINE, SCOPUS, EMBASE, INDEX COPERNICUS,
MINISTRY OF SCIENCE AND HIGHER EDUCATION, POLISH MEDICAL BIBLIOGRAPHY

Copyright: © ALUNA Publishing

Articles published on-line and available in open access are published under Creative Commons Attribution-Non Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.

Distribution and Subscriptions:

Bartosz Guterman prenumerata@wydawnictwo-aluna.pl

Graphic design / production:

Grzegorz Sztank

fajne.work

Publisher:

ALUNA Publishing
29 Przesmyckiego st.,
05-510 Konstancin – Jeziorna, Poland
www.wydawnictwo-aluna.pl
www.wiadomoscilekarskie.pl
www.wiadlek.pl

Wiadomości Lekarskie

Medical Advances

Official journal of the Polish Medical Association
Wiadomości Lekarskie has been published since 1928



Editorial Team

Editor in-Chief:

Prof. Paweł Kalinski – Buffalo, USA

Honorary Editor in-Chief:

Prof. Władysław Pierzchała – Katowice, Poland

Deputy Editor in-Chief:

Prof. Waldemar Kostewicz – Warsaw, Poland
President Polish Medical Association

Statistical Editor:

Dr Lesia Rudenko – Konstancin-Jeziorna, Poland

Managing Editor:

Agnieszka Rosa – amarosa@wp.pl

International Editorial Office:

Nina Radchenko (editor) – n.radchenko@wydawnictwo-aluna.pl

International Editorial Board – in-Chief:

Marek Rudnicki Chicago, USA

International Editorial Board – Members:

Stalbek M. Akhunbaev	Bishkek, Kyrgyzstan	Jerzy Robert Ładny	Białystok, Poland
Kris Bankiewicz	San Francisco, USA	Stella Nowicki	Memphis, USA
Christopher Bara	Hannover, Germany	Alfred Patyk	Gottingen, Germany
Krzysztof Bielecki	Warsaw, Poland	Palmira Petrova	Yakutsk, Russia
Zana Bumbuliene	Vilnius, Lithuania	Waldemar Priebe	Houston, USA
Stanislav Czudek	Ostrava, Czech Republic	Maria Siemionow	Chicago, USA
Mowafaq Muhammad Ghareeb	Baghdad, Iraq	Aleksander Sieroń	Katowice, Poland
Nataliya Gutorova	Kharkiv, Ukraine	Vladyslav Smiianov	Sumy, Ukraine
Marek Hartleb	Katowice, Poland	Tomasz Szczepański	Katowice, Poland
Roman Jaeschke	Hamilton, Canada	Andrzej Witek	Katowice, Poland
Andrzej Jakubowiak	Chicago, USA	Jerzy Woy-Wojciechowski	Warsaw, Poland
Peter Konturek	Saalfeld, Germany	Zbigniew Wszolek	Jacksonville, USA
George Krol	New York, USA	Vyacheslav Zhdan	Poltava, Ukraine

 CONTENTS

ORIGINAL ARTICLES

Yuliia Cherniavska, Alina Davydenko, Valerii Pokhylko, Liliia Fishchuk, Zoia Rossokha

- Preliminary study of the influence of maternal and neonatal NOS3 (rs1799983), IL1B (rs1143634) genes variants and their intergenic interaction on the development of hypoxic-ischemic encephalopathy in newborns in the context of treatment planning** 2373

Randa Saad Hameed, Rihab Hameed Almudhafar, Liwaa Hussein Mahdi

- Immunohistochemical evaluation of GATA-3 in patients with urinary bladder cancer** 2381

Ivan Mamontov, Tamara Tamm, Kostiantyn Kramarenko, Dmytro Ryabushchenko, Dmytro Sytnik, Samer Dghaili

- Small common bile duct – the risk factor for post-ercp pancreatitis in patients with choledocholithiasis** 2388

Olena Vysotska, Olena Falova, Liubov Rysovana, Oksana Platyniuk, Oleksandr Bazhenov

- Woman's self-realization in the system and structure of family relations** 2394

Maryana M. Shevchuk, Andriana P. Skalat, Khristina B. Burda, Nataliya S. Magera, Olesia I. Martovlos

- Evaluation of the clinical effectiveness of conservative (pre-surgical) treatment schemes for generalized periodontitis in patients with various somatic pathologies** 2403

Vladyslav Tuliakov, Oleksander Barkov

- Biomarkers of inflammation in the blood of patients with the degenerative lumbar spine disease and complications after transpedicular fixation** 2412

Amina Jabri, Zaid Fakhruddin, Evan Hameed

- Assessment of prescribing practice of anti-hypertensive medications in a sample of iraqi patients with diabetes** 2419

Yevhen Mykhaliuk, Yehor Horokhovskiy, Anatolii Bosenko, Oleg Bazylchuk, Mykhailo Khoroshukha, Nadiia Orlyk, Mariia Topchii

- Heart rate and blood pressure in soccer players differing in sports qualification** 2426

Yurii O. Hrubar, Iryna Ya. Hrubar, Nadiia M. Hrabyk, Markiiian Yu. Grubar, Yuliana Yu. Hrubar

- Change of the anterior talus-fibular ligament length of asymptomatic patients during inversion stress test** 2435

Lubna Sami Ali, Ali M. Janabi

- Phytochemical analysis of ethanolic leaf extract of Cordia Myxa and its anti-inflammatory and cytotoxic activities** 2442

Radostina Hristova Simeonova, Ignat Ignatov, Zlatin Ivanov, Georgi Nanov

- Radial shock wave therapy with "Intellect PRW Lite" for calcifying tendinopathy of the shoulder, comparative effectiveness with conventional physical therapy and follow up data** 2451

Change of the anterior talus-fibular ligament length of asymptomatic patients during inversion stress test

Yurii O. Hrubar¹, Iryna Ya. Hrubar², Nadiia M. Hrabyk², Markiian Yu. Grubar³, Yuliana Yu. Hrubar³

¹TERNOPIL I. IA. HORBACHAVSKYI NATIONAL MEDICAL UNIVERSITY, TERNOPIL, UKRAINE

²TERNOPIL VOLODYMYR HNATIUK NATIONAL PEDAGOGICAL UNIVERSITY, TERNOPIL, UKRAINE

³MUNICIPAL NONPROFIT ENTERPRISE «TERNOPIL REGIONAL CLINICAL HOSPITAL», TERNOPIL REGIONAL RADA, TERNOPIL, UKRAINE

ABSTRACT

Aim: To study the ATFL length among asymptomatic population of young patients on both joints by using the method of ultrasound examination of the ankle joint and applying the inversion stress test.

Materials and Methods: the length of the ATFL was measured among 60 boys and 60 girls during the research. Sonography was performed on both joints, in each of the two studied positions, three times per position in order to obtain the average ligament length. A total of 720 examinations were conducted.

Results: According to the results of sonographic research, the average length of the ATFL of the right ankle joint of young men in a neutral position was $19,08 \pm 0,3$ mm, of the left one – $19,08 \pm 0,31$ mm. During the inversion stress test, the length of the ligament increased on the right up to $19,65 \pm 0,3$ mm, on the left – up to $19,63 \pm 0,3$ mm. The average length of the ATFL of the right ankle joint of girls in neutral position was $16,92 \pm 0,2$ mm, of the left one – $16,9 \pm 0,19$ mm. During the inversion stress test, the length of the ligament increased on the right up to $17,37 \pm 0,2$ mm, on the left – up to $17,36 \pm 0,20$ mm. Statistical difference of the length of the ATFL of both joints between the neutral position and the stress test was at the level $p < 0,001$ among boys and girls.

Conclusions: Comparison of the length of the anterior talus-fibular ligament of boys and girls in different positions during testing of both joints indicates a statistically significant difference at the level ($P < 0,001$).

KEY WORDS: ankle joint, anterior talus-fibular ligament, inversion stress test, stress sonography

Wiad Lek. 2024;77(12):2435-2441. doi: 10.36740/WLek/195548 DOI

INTRODUCTION

Trauma of ligaments of the ankle joint belongs to the most common injuries, both in everyday life and in sports. It is the reason for requests for urgent medical help in 20% of cases, however, total number of these injuries is much higher, since less than half of all people with ligament damage of this specified localization seek qualified medical assistance [1].

Patients with ligament damage are usually young and physically active people between 14 and 24 years old. [2]. More than 65% of cases are isolated injuries of the anterior talus-fibular ligament (ATFL). Interest in this ligament is due to the fact that it is considered the main means of protection against injuries, associated with ankle joint inversion combined with plantar flexion [3].

The consequences of ligament damage can lead to a number of chronic residual symptoms, which are diagnosed among from 40% to 50% of patients and include injury recurrence, chronic pain, chronic instability, osteoarthritis, adverse psychological effects [4].

The diagnosis of ligament damage is established on the basis of the patient's complaints, the mechanism of trauma occurrence, study of the local status, conducting of diagnostic tests and use of additional diagnostic methods. It is well known that there is limited evidence of the use of many specific orthopedic tests, which are widely used at injuries of ankle joint ligaments [5].

Among the non-invasive instrumental methods of diagnosis, they most often use stress radiography and magnetic resonance tomography. Stress radiography is often not informative enough to establish a final diagnosis [6]. Magnetic resonance imaging is the most accurate diagnostic method for assessing ligament damage, considering its high resolution and accuracy, however, its use in case of an acute injury of the specified location is not considered appropriate [7].

In order to improve the results of diagnostics, sophisticated devices and structures are introduced, but they are unlikely to be widely used in everyday practice during study of the state of the ATFL [8, 9].

Today, sonography is becoming increasingly important for the assessment of the ligamentous system of the ankle joint because of low cost, speed, availability and absence of ionizing radiation [10]. Unlike MR- images, it allows to perform dynamic maneuvers, which increases the level of visualization of intact ligaments and improves localization of their injuries [11-13].

Dynamic sonography has a high potential for detecting ankle joint ligament instability, but the parameters of the normative values of change of the length of the anterior talus-fibular ligament during stress tests in different age groups have not been fully determined.

AIM

To study changes of the anterior talus-fibular ligament length among asymptomatic population of young patients by using the device and the method of ultrasound examination of the ankle joint during applying the inversion stress test on both ankle joints.

MATERIALS AND METHODS

The research was conducted using an ultrasound console Acuson Antares (Siemens) applying a high-frequency broadband linear sensor with the operating frequency 7–12 MHz and using a standard acoustic gel.

The researcher visualized and measured the length of the anterior talus-fibular ligament of all examined persons in two positions, three times per position, in order to obtain the average length of the ligament. The examined persons relaxed the lower limb in the intervals between measurements, in order to avoid possible distortions of motion range due to the development of muscle contracture.

In the first position (position A), the examined persons were lying on the couch, the examined lower limb was placed on the device for ultrasound examination of the ankle joint in the position of bending in the knee joint at an angle 45°, and the area of the ankle joint hung freely in the position of plantar flexion at the angle 20° (Fig. 1. A.). This position was defined as the neutral resting position and as the baseline value for each subsequent measurement of the ATFL length.

The sonographic sensor was placed in accordance with the anatomical localization of the ATFL almost parallel to the sole of the foot (Fig. 1. B.)

A. Neutral position of the examined lower limb, the lower leg is placed on the ultrasound device. B. Location of the sonographic sensor for sonography of the anterior talus-fibular ligament in a neutral position.

During the sonographic examination, the distance from the anterolateral edge of the lateral bone was measured, which corresponds to the beginning of the ATFL, to the anterolateral corner of the lateral surface of the talus, which corresponds to the anatomical location of ligament fixation. The length of the anterior talus-fibular ligament was determined as a straight line between these two bony reference points, and was measured on a longitudinal ultrasound image made along the course of its fibers (Fig. 2. A, B).

In the second position (position B, of maximum inversion): examined persons were lying on the examination table, in the standard starting position. The researcher performed maximum inversion in the ankle joint till tissue resistance began.

The length of the anterior talus-fibular ligament in the position of the inversion stress test was defined as a straight line between the bone reference points described above, and measured on a longitudinal ultrasound image, which was performed along the course of its fibers (Fig. 3. A, B).

STATISTICAL ANALYSIS

Statistical analysis was performed using the software Statistica 8.0. Results were presented as average values with 95% confidence intervals. The Shapiro-Wilk test was used to confirm whether the data were normally distributed. Student's criterion was used to compare the continuous data. A Wilcoxon signed-rank test was performed to compare ultrasound results for the right and the left ankle.

RESULTS

The research was conducted in 2022-2023. The students of Ternopil National Medical University voluntarily participated in it. We examined the ankle joints of both limbs of 60 boys and 60 girls. The sample did not include persons with a medical history including injury of ligament of the ankle joint, operative interventions of the specified area, positive tests for joint hypermobility syndrome. All enrolled subjects signed a consent form prior to participating in the study.

Sonography was performed three times in a neutral position and during the inversion stress test. The average length of the anterior talus-fibular ligament of the right and the left ankle joint in each position was included in the protocol. In total 720 tests were carried out.

The morphometric characteristics of the examined persons is given in the Table 1.

According to the results of sonographic research, average length of the anterior talus-fibular ligament



Fig. 1. Neutral position for examining the length of the anterior talus-fibular ligament.

Table 1. Morphometric characteristics of the examined persons

Parameters	Boys	Girls
Age, years	22,63±0,11	22,54±0,12
Height, cm	181,42±0,74	168,68±0,69
Foot length, size cm	42,87±0,15	37,97±0,16
Body weight, kg	76,38±1,22	59,63±0,92
Body weight index, conventional unit	21,14±0,28	18,05±0,25

Table 2. The length of the anterior talus-fibular ligament of the ankle joint of the examined persons in different test positions

Group	Test state	The length of the anterior talus-fibular ligament, the right ankle joint	The length of the anterior talus-fibular ligament, the left ankle joint	Statistical difference between the ligaments of the right and the left joint, P	Statistical difference of ligament length in different positions, the right joint, P	Statistical difference of ligament length in different positions, the left joint, P
Boys	Neutral position, mm	19,08±0,3	19,08±0,31	>0,05	<0,001	<0,001
	Stress test, mm	19,65±0,3	19,63±0,3	>0,05		
	Difference of ligament length, mm	0,57	0,56	>0,05		
Girls	Neutral position, mm	16,92±0,2	16,9±0,19	>0,05	<0,001	<0,001
	Stress test, mm	17,37±0,2	17,36±0,2	>0,05		
	Difference of ligament length, mm	0,45	0,45	>0,05		

of the right ankle joint of young men in a neutral position was 19,08±0,3 mm (Table II). The length of

the ligament increased up to 19,65±0,3 mm during maximum inversion stress test. The difference of




Fig. 2. Length measurement of the anterior talus-fibular ligament (ATFL) in a neutral position during sonography.

A. Dimensions of the anterior talus-fibular ligament in a neutral position of the examined person, male (A-B measurement points of the anterior talus-fibular ligament in a neutral position, 1 – splintbone; 2 – talus).

B. Dimensions of the anterior talus-fibular ligament in a neutral position of the examined person, female (A-B measurement points of the anterior talus-fibular ligament in a neutral position, 1 – splintbone; 2 – talus).




Fig. 3. Length measurement of the anterior talus-fibular ligament in the position of the inversion stress test during sonography.

A. Dimensions of the anterior talus-fibular ligament in a maximum position of the inversion stress test of the examined person, male (A-B measurement points of the anterior talus-fibular ligament, 1 – splintbone; 2 – talus).

B. Dimensions of the anterior talus-fibular ligament in a maximum position of the inversion stress test of the examined person, female (A-B measurement points of the anterior talus-fibular ligament, 1 – splintbone; 2 – talus).

the ligament length between neutral position and maximum inversion was 0,57 mm. 95% DI in the neutral position was 0,078 mm (19,01-19,16 mm), and during the inversion stress test – 0,078 mm (19,58-19,73 mm). ATFL extension in percentage terms was 2,99%. Statistical difference of the length of the anterior talus-fibular ligament of the right ankle joint between the neutral position and the stress test was at the level of $p < 0,001$.

As for girls, average length of the anterior talus-fibular ligament of the right ankle joint in a neutral position was $16,92 \pm 0,2$ mm (Table 2). At maximum inversion it increased up to $17,37 \pm 0,2$ mm. 95% DI amounts 0,05 mm in both positions: in a neutral position within 16,87-16,97 mm, and during the stress test – 17,32-17,42 mm. The difference of the ligament length between neutral position and maximum inversion was 0,45 mm. In

percentage terms it was 2,67%. As for girls, statistical difference of the length of the anterior talus-fibular ligament of the right ankle joint between the neutral position and the stress test was at the level of $p < 0,001$.

Sonographic research of the ATFL the left ankle joint of girls showed that in the neutral position its length was $16,9 \pm 0,19$ mm, and during the inversion stress test – $17,36 \pm 0,20$ mm. 95% DI (confidence interval) in both positions was 0.05 mm: in neutral position it was within the limits 16,85-16,95 mm, and during the inversion stress test – 17,31-17,42 mm. The difference of the ligament length between neutral position and maximum inversion was 0,45 mm, and in percentage terms it was 2,66%. As for girls, statistical difference of the length of the anterior talus-fibular ligament of the left ankle joint between the neutral position and the stress test was at the level of $p < 0,001$.

Our research shows that statistically significant difference between the parameters of the ATFL ligament of the right and the left ankle joint is absent in the group of boys and girls, both in the neutral position and during the inversion stress test ($P>0,05$).

Length comparison of the ATFL of boys and girls in different positions during testing of both joints indicates a statistically significant difference between the parameters of boys and girls at the level of $p<0,001$.

Difference of the ligament length of the right ankle joint of boys is 0,57 mm, and of girls – 0,45 mm. A similar situation is observed with the length of the anterior talus-fibular ligament of the left ankle joint in different positions: 0,56 and 0,45 mm.

DISCUSSION

With increasing the evidence of efficiency and accuracy of sonography of the musculoskeletal system, more and more attention is paid to ultrasound methods of diagnosing injuries of the anterior talus-fibular ligament (ATFL). So, Oae K. et. al. Informed about the high accuracy of using longitudinal ultrasound examination of ATFL at identification of morphological changes indicating its damage [14]. Study of ligament dimensions was carried out both on cadaveric material and by instrumental methods. Inchai C. et al., while studying fresh-frozen cadavers with ages at death ranging from 25 to 90 years old [15] established, that average dimensions of the anterior talus-fibular ligament were $19,54\pm 2,96$ mm. During MRI studies, the average length of ATFL was from $21,5\pm 0,5$ mm to $24,5\pm 3,3$ mm and significant differences in size between its one- and two-bundle structure were not found [16].

Longitudinal sonographic measurement between bony attachment ATFL showed that its visible length varies in the neutral position from $16,1\pm 3,63$ mm to $21,73\pm 2,67$ mm [17]. With inversion load, it can increase up to $23,14\pm 2,49$ mm [18].

The results of our research show that the average length of the right of the ATFL of boys in the neutral position was $19,08\pm 0,3$ mm. Average length values increased to $19,65\pm 0,3$ mm during the maximum inversion stress test. Average length of the left anterior

talus-fibular ligament of boys in the neutral position was $19,08\pm 0,31$ mm. It increased to $19,63\pm 0,3$ mm during maximum inversion. Average values of the length of both the right and the left anterior talus-fibular ligament of girls in the neutral position were $16,92\pm 0,2$ mm for the right ankle joint and $16,9\pm 0,19$ mm for the left ankle joint. It increases to $17,37\pm 0,2$ mm to $17,36\pm 0,2$ mm accordingly during the inversion stress test. Thus, the average length of the ATFL of both boys and girls according to the results of our research corresponds with the data obtained by Mizrahi D.J et. al. [19]. Extension of the ATFL of boys in percentage terms it was 2,99% for the right ankle joint and 2,88% for the left ankle joint. Extension of the ligament of girls accordingly was 2,67% and 2,66%. These results correspond with the data obtained by Jeys L. et al. [20], which indicate that change of the ATFL length during movement is less than 5%.

The results of our research regarding the definition of the average length of ATFL of boys and girls in neutral position and during inversion are within the limits reported by Croy et al. for neutral position and inversion, $18,6\pm 1,5$ mm and $19,9\pm 2,3$ mm respectively [21]. The length of the anterior talus-fibular ligament both in the neutral position and during maximum inversion stress test of boys compared to girls was longer ($P<0,001$).

Potential limitation of this research was that the exact force applied during manual loading wasn't measured, however, maximal loading in plantar flexion and talar bone inversion can be reproduced in clinical conditions.

CONCLUSIONS

Our research shows that statistically significant difference between the parameters of the ATFL ligament of the right and the left ankle joint is absent in the group of boys and girls, both in the neutral position and during the inversion stress test ($P>0,05$).

Length comparison of the ATFL of boys and girls in different positions during testing of both joints indicates a statistically significant difference between the parameters of boys and girls at the level of $P<0,001$.

REFERENCES

1. Cavazos GJJr, Harkless LB Harkless LB. The epidemiology, evaluation, and assessment of lateral ankle sprains in athletes. *Journal of Sports Medicine and Therapy*. 2021;6:8–17. doi:10.29328/journal.jsmt.
2. Martin R, Davenport T, Paulseth S et al. Ankle Stability and Movement Coordination Impairments: Ankle Ligament Sprains Clinical Practice Guidelines. *J Orthop Sports Phys Ther*. 2013;43(9):A1-40. doi: 10.2519/jospt.2013.0305.
3. Dalmau-Pastor M, El-Daou H, Stephen JM. Clinical Relevance and Function of Anterior Talofibular Ligament Superior and Inferior Fascicles: A Robotic Study. *Am J Sports Med*. 2023;51(8):2169-2175. doi: 10.1177/03635465231172196.

4. Aicale R, Maffulli N. Chronic Lateral Ankle Instability: Topical Review. *Foot Ankle Int.* 2020;41(12):1571-1581. doi: 10.1177/1071100720962803.
5. Larkins LW, Baker RT, Baker JG. Physical Examination of the Ankle: A Review of the Original Orthopedic Special Test Description and Scientific Validity of Common Tests for Ankle Examination. *Arch Rehabil Res Clin Transl.* 2020;2(3):100072. doi: 10.1016/j.arrct.2020.100072.
6. Cao M, Liu S, Zhang X et al. Imaging diagnosis for anterior talofibular ligament injury: a systemic review with meta-analysis. *Acta Radiol.* 2023;64(2):612-624. doi: 10.1177/02841851221080556.
7. Dalal S, Morgan G. Chronic lateral ankle instability: Results of two-staged approach, correlation with magnetic resonance imaging findings, and incidence of associated pathologies – A 4-year follow-up study. *J Arthrosc Surg Sport Med* 2020;1(2):178-85. doi:10.25259/JASSM_34_2020.
8. Wenning M, Gehring D, Lange T et al. Clinical evaluation of manual stress testing, stress ultrasound and 3D stress MRI in chronic mechanical ankle instability. *BMC Musculoskelet Disord.* 2021;22(1):198. doi: 10.1186/s12891-021-03998-z.
9. Teramoto A, Murahashi Y, Shoji H. Quantitative Evaluation of Ankle Instability Using a Capacitance-Type Strain Sensor. *Foot Ankle Int.* 2021;42(8):1074-1080. doi: 10.1177/1071100721996714.
10. Ekinci S, Polat O, Günalp M et al. The accuracy of ultrasound evaluation in foot and ankle trauma. *Am J Emerg Med.* 2013;31(11):1551-5. doi: 10.1016/j.ajem.2013.06.008.
11. Baloch N, Hasan OH, Jessar M et al. "Sports Ultrasound", advantages, indications and limitations in upper and lower limbs musculoskeletal disorders. Review article. *Int J Surg.* 2018;54(B):333-340. doi: 10.1016/j.ijssu.2017.11.034.
12. Hrubar YuO, Hrubar MYu, Koptiyukh VV. «Prystryi dlya ul'trazvukovoho doslidzhennya homilkovostopnoho suhloba» Patent na korysnu model' № 138492 Ukrayina, MPK A61B 8/00; (2019.01); Zayavleno 30.05.2019; Opublikovano 25.12.2019. Byuleten' №22. [«Device for ultrasonic research of the hoof moisture» Utility model patent No. 138492 of Ukraine, MPK A61B 8/00; (2019.01); Filed on 05/30/2019; Published on 12/25/2019. Bulletin No. 22]. (Ukrainian)
13. Hrubar YuO, Hrubar MYu, Koptiyukh VV. «Sposib ul'trazvukovoho doslidzhennya homilkovostopnoho suhloba» Patent na korysnu model' № 138901 Ukrayina, MPK A61B 8/00; (2019.01); Zayavleno 30.05.2019; Opublikovano 10.12.2019. Byuleten' №23. [“Method of ultrasonic examination of the homilial dryness” Utility model patent No. 138901 Ukraine, MPK A61B 8/00; (2019.01); Filed 30.05.2019; Published 10.12.2019. Bulletin No. 23]. (Ukrainian)
14. Oae K, Takao M, Uchio Y et al. Evaluation of anterior talofibular ligament injury with stress radiography, ultrasonography and MR imaging. *Skeletal Radiol.* 2010;39(1):41-7. doi: 10.1007/s00256-009-0767-x.
15. Inchai C, Vaseenon T, Tanaka Y et al. The Specific Anatomical Morphology of Lateral Ankle Ligament: Qualitative and Quantitative Cadaveric based Study. *Orthop Sur.* 2023;15:2683–2688. doi:10.1111/os.13872.
16. Szaro P, Gataa KG, Solidakis N et al. Morphometric relationships between dimensions the anterior talofibular ligament and calcaneofibular ligament in routine magnetic resonance imaging. *J Exp Orthop.* 2021;8:90. doi: 10.1186/s40634-021-00406-22021.
17. Kristen KH, Seilern J, Wiedemann J et al. Reliability of ultrasonography measurement of the anterior talofibular ligament (ATFL) length in healthy subjects (in vivo), based on examiner experience and patient positioning. *J Exp Orthop.* 2019;6(1):30. doi: 10.1186/s40634-019-0199-z.
18. Kikumoto T, Akatsuka K, Nakamura E et al. Quantitative evaluation method for clarifying ankle plantar flexion angles using anterior drawer and inversion stress tests: a cross-sectional study. *J Foot Ankle Res.* 2019;12:27. doi: 10.1186/s13047-019-0337-y.
19. Mizrahi DJ, Nazarian LN, Parker L. Evaluation of the Anterior Talofibular Ligament via Stress Sonography in Asymptomatic and Symptomatic Populations. *J. Ultrasound Med.* 2018;37(8):1957-1963. doi: 10.1002/jum.14542.
20. Jeys L, Korrosis S, Stewart T et al. Bone anchors or interference screws? A biomechanical evaluation for autograft ankle stabilization. *Am J Sports Med.* 2004;32(7):1651-9. doi: 10.1177/0363546504265051.
21. Croy T, Saliba S, Saliba E et al. Differences in lateral ankle laxity measured via stress ultrasonography in individuals with chronic ankle instability, ankle sprain copers and healthy individuals. *J Orthop Sports Phys Ther.* 2012;42(7):593-600. doi: 10.2519/jospt.2012.3923.

CONFLICT OF INTEREST

The Authors declare no conflict of interest

CORRESPONDING AUTHOR

Yurij O. Hrubar

Ternopil National Medical University
1 Maidan Voli, 46001 Ternopil, Ukraine
e-mail: Yuhrubar@gmail.com

ORCID AND CONTRIBUTIONSHIP

Yurii O. Hrubar: 0000-0002-4221-2250 **B** **C** **D**

Iryna Ya. Hrubar: 0000-0002-0809-1299 **A** **D** **E**

Nadiia M. Hrabyk: 0000-0002-8882-9782 **C** **D** **F**

Markiiian Yu. Grubar: 0000-0002-4696-0213 **B** **D** **E**

Yuliana Yu. Hrubar: 0000-0003-0951-9485 **A** **D** **F**

A – Work concept and design, **B** – Data collection and analysis, **C** – Responsibility for statistical analysis, **D** – Writing the article, **E** – Critical review, **F** – Final approval of the article

RECEIVED: 28.03.2024

ACCEPTED: 04.11.2024